

Equisedan[®] Vet

Registered pursuant of the ACVM Act 1997 A10285

Detomidine HCL 10mg/mL

**Manufactured to European Union [EU] Good Manufacturing Practice [GMP].
A guarantee of product quality and reliability.**

- ✓ Registered and introduced in a number of European Community [EC] countries.¹
- ✓ Superior antinociceptive (reduction in pain sensitivity) compared with xylazine or romfidine in horses.²
- ✓ Detomidine + butorphanol demonstrated “greater and longer muscle relaxation” and “confirmed detomidine + butorphanol combination is safer and more appropriate for painful procedures in the standing horse”.³
- ✓ Safe, effective sedation was demonstrated when detomidine was administered immediately following intensive exercise in thoroughbreds.⁴
- ✓ For colic, detomidine provided significantly more analgesia than flunixin or xylazine for abdominal pain in clinical cases of colic.⁵
- ✓ A recent study reported that multiple administrations of detomidine in the last trimester of pregnancy in mares had no measurable effects on the outcome of the pregnancy.⁶

Presentation: 10mg/mL detomidine HCL – 10mL multi-dose vial.

Equisedan Vet

1. Data on File

2. Moens Y *et al* Vet. Anaesthesia and Analgesia, 30 (3), 2003

3. Kruljc P *et al* Acta Veterinaria Hungarica, 54 (1) 2006

4. Hubbel JA *et al* Am J Vet. Research, 60 (10), 1999

5. Jochel W *et al* Equine Vet. Supp. Jun, (7) 1989

6. Luukanen L *et al* E Vet J, 29 (5) 1997

ausrichter

www.ausrichter.com

info@ausrichter.com

Predictable, dose-related sedation and analgesia in horses

Equisedan[®] Vet

10mg/mL Solution for Injection

Detomidine HCL 10mg/mL

**Manufactured to European Union [EU] Good Manufacturing Practice [GMP].
A guarantee of product quality and reliability.**

- ▶ Registered and introduced in a number of European Community [EC] countries.¹
- ▶ Superior antinociceptive (reduction in pain sensitivity) compared with xylazine or romfidine in horses.²
- ▶ Detomidine + butorphanol demonstrated "greater and longer muscle relaxation" and "confirmed detomidine + butorphanol combination is safer and more appropriate for painful procedures in the standing horse".³
- ▶ Safe, effective sedation was demonstrated when detomidine was administered immediately following intensive exercise in thoroughbreds.⁴
- ▶ For colic, detomidine provided significantly more analgesia than flunixin or xylazine for abdominal pain in clinical cases of colic.⁵
- ▶ A recent study reported that multiple administrations of detomidine in the last trimester of pregnancy in mares "had no measurable effects on the outcome of the pregnancy."⁶

Presentation: 10mg/mL detomidine HCL – 10mL multi-dose vial.

1. Data on File

2. Moens Y *et al*: Vet. Anaesthesia and Analgesia, 30 (3), 2003

3. Kruljic P *et al*: Acta Veterinaria Hungarica, 54 (1) 2006

4. Hubbel JA *et al*: Am J Vet. Research, 60 (10), 1999

5. Jochel W *et al*: Equine Vet. Supp. Jun, (7) 1989

6. Luukanen L *et al*: E Vet J, 29 (5) 1997

ausrichter

Ausrichter Pty Ltd

ABN 79 000 908 529

2/21 Chester Street, Camperdown NSW 2050

Telephone: (02) 9517 1166 Fax: (02) 9516 5810

Email: info@ausrichter.com www.ausrichter.com

Animal Health Products

APVMA Approval No.: 61880/0109

Manufactured by

richter pharma ag

Wels, Austria



European Union